

Peidiwch â buddsoddi mewn afiechyd

Mae symud stoc yn creu risg o drosglwyddo afiechyd bob amser. Mae anifeiliaid a brynwyd yn risg i'ch buches, ac mae anifeiliaid a brynwyd hefyd mewn perygl o godi afiechyd o'r anifeiliaid yn eich buches. Bydd cyflwyno afiechyd yn arwain at ganlyniadau niweidiol i iechyd, lles, cynhyrchiant a phroffidioldeb.



TWBERCWLOSIS GWARTHEG

Gall camau i gadw anifeiliaid ar wahân wedi'u cyfuno â phrofi anifeiliaid yn breifat ar ôl eu symud leihau'r perygl o gael colledion yn y tymor hir a byr.

Canfyddwch beth yw hanes profi TB y fferm ac o'r ardal yr ydych yn prynu ohoni.

Diogelwch siediau gwartheg, ardaloedd bwydo a storfeydd rhag fywyd gwyllt.

Os yw'r gwartheg yn cael eu bwydo allan ar borfa, canfyddwch ffyrdd i ddiogelu bwyd a chafnau oddi wrth bywyd gwyllt.

Ystyriwch ddewisiadau eraill yn hytrach na blociau pan allan ar borfa.

Sicrhewch bod perimedr y fferm yn atal cyswllt gyda da byw drws nesaf.

Ffensiwch setiau a geudai moch daear i ffwrdd ble bynnag posib.



PERYGLON PARASITIAID

LLYNGYR Y PERFEDD
LLYNGYR YR AFU
LLYNGYR YR YSGYFAINT
ECTOPARASITIAID

Sicrhau bod triniaethau yn cael eu defnyddio yn gywir i fod yn effeithiol ac er mwyn atal ymwrthedd

Gwelwch gwefan: www.cattleparasites.org.uk am wybodaeth ar barasitiaid gwartheg a'u rheolaeth.

Adolygwch eich Rhaglen Rheoli Parasitiaid gyda help eich milfeddyg (cynllunio iechyd).

Llyngyr yr Ysgyfaint - gwiriwch statws brechu.

Ectoparasitiaid (trogod, llau, mansh) - archwiliwch wartheg yn ofalus a'u trin pan fydd angen.



OSGOI CLOFFNI

DERMATITIS DIGIDOL
DRWGYN Y DROED

Mae'r rhain yn afiechydon heintus!

Mae Dermatitis Digidol yn afiechyd heintus difrifol.

Gwiriwch y drefn trin a hanes buches y gwerthwr.

Defnyddiwch fath traed penodol ar gyfer stoc newydd fel y cytunwyd yn eich cynllun iechyd anifeiliaid gyda'ch cyngorydd milfeddygol.



AFIECHYDON CUDD

BVD
IBR
LEPTOSBIROSIS (MILHEINTIAU)
NEOSPOR
JOHNES

Gall wartheg sydd ddim yn dangos afiechyd fod yn "gludwyr" a gallent fod yn lledaenu heintiau o fewn y fuches!

Prynwch oddi wrth ffynonellau yr ydych yn ymddiried ynddyn nhw.

Prynwch stoc achrededig.

Ymchwiliwch i unrhyw dystiolaeth o afiechyd.

Buddsoddwch ar gyfer buddion tymor byr, canolig a hir er mwyn helpu i ddiogelu eich diwydiant drwy ddechrau rhaglen rheoli afiechyd neu achrediad rhydd o glefyd.

- Mae gwybod statws afiechyd eich buches yn hanfodol.
- Prynwch o ffynonellau yr ydych yn gwybod eu statws iechyd.
- Mabwysiadwch fesurau cadw anifeiliaid ar wahân fel y nodir yn eich cynllun iechyd anifeiliaid.
- Ymchwiliwch i unrhyw arwyddion o afiechyd.
- Gofynnwch am gyngor eich milfeddyg.

Gall afiechyd heintus ledaenu trwy esgidiau, dillad ac offer! Os defnyddir bath traed gyda diheintydd, sicrhewch fod y diheintydd yn gywir gyda'r gwanhad cywir. Mabwysiadwch lefelau uchel o fioddiogelwch wrth ymweld â ffermydd eraill, marchnadoedd neu ladd-dai.

Don't invest in disease

Movement of stock always carries the risk of disease transfer. Purchased animals pose a risk to your herd, and purchased animals are also at risk of picking up disease from the animals in your herd. The introduction of disease will have adverse consequences for health, welfare, productivity and profitability.

- Knowing the disease status of your own herd is crucial
- Only buy from sources of known health status
- Adopt quarantine measures as specified in your animal health plan
- Investigate any signs of disease
- Seek guidance from your vet

Infectious disease can spread via boots, clothing and equipment! If using a disinfectant foot dip ensure the correct disinfectant at the correct dilution. Adopt high levels of biosecurity when visiting other farms, markets or abattoirs.



BOVINE TUBERCULOSIS

Quarantine measures combined with private post-movement testing can reduce the danger of losses both in the long and short term.

Know the TB testing history of the farm and of the locality from where you are purchasing stock.

Protect your cattle housing and feed stores from wildlife.

If feeding at pasture, find ways of protecting feed and troughs from wildlife.

Consider alternatives to supplementary feed blocks at pasture.

Ensure that the farm perimeter prevents contact with neighbouring livestock.

Wherever possible, fence off badger setts and latrines.



PARASITE PERIL

INTESTINAL WORMS
LIVER FLUKE
LUNG WORM
ECTOPARASITES

Ensure treatments are used correctly to be effective and to prevent resistance.

See: www.cattleparasites.org.uk for information on cattle parasites and their control.

Regularly review your Parasite Control Programme with help from your vet (health planning).

Lung Worm – check vaccination status.

Ectoparasites (ticks, lice, mange) - examine cattle carefully and treat when necessary.



LAME EXCUSES?

DIGITAL DERMATITIS
FOUL IN THE FOOT

These are infectious diseases!

Digital Dermatitis is a major infectious disease.

Check treatment regime and herd history of vendor.

Use a specific footbath for new arrivals as agreed in your animal health plan with your veterinary adviser.



TICKING TIMEBOMBS

BVD
IBR
LEPTOSPIROSIS (ZONOSSES)
NEOSPORA
JOHNES


Cattle showing no ill-health can be “carriers” and spread infection within the herd!

Buy from trusted sources.

Buy accredited stock.

Investigate evidence of disease.

Invest for short, medium and long term benefits and help safeguard your industry by starting a disease control program or disease freedom accreditation.



www.menterabuses.co.uk/cyswlltffermio
www.menterabuses.co.uk/farmingconnect



Llywodraeth Cymru
Welsh Government



Cronfa Amaethyddol Ewrop ar gyfer
Datblygu Gwledig: Ewrop yn Buddsoddi
mewn Ardalwedd Gwledig
The European Agricultural
Fund for Rural Development:
Europe Investing in Rural Areas